

**ENVIRONMENTAL POLLUTION, PESTICIDES, AND THE PREVENTION OF CANCER:
MISCONCEPTIONS***

Bruce N. Ames and Lois Swirsky Gold

Dr. Bruce N. Ames is a Professor of Biochemistry and Molecular Biology and Director of the National Institute of Environmental Health Sciences Center, University of California, Berkeley, CA 94720. He is a member of the National Academy of Sciences and was on their Commission on Life Sciences. He was a member of the National Cancer Advisory Board of the National Cancer Institute (1976-82). His many awards include: the General Motors Cancer Research Foundation Prize (1983), the Tyler Prize for environmental achievement (1985), the Gold Medal Award of the American Institute of Chemists (1991), the Glenn Foundation Award of the Gerontological Society of America (1992), the Lovelace Institutes Award for Excellence in Environmental Health Research (1995), the Honda Foundation Prize for Ecotoxicology (1996) and the Japan Prize (1997). His 380 publications have resulted in his being the 23rd most-cited scientist (in all fields) (1973-84). BNAmes@UCLink4.Berkeley.edu (510)642-5165

Dr. Lois Swirsky Gold is Director of the Carcinogenic Potency Project at the National Institute of Environmental Health Sciences Center, University of California, Berkeley, CA 94720 and a Senior Scientist at the Berkeley National Laboratory, Berkeley, CA 94720. She has published 85 papers on analyses of animal cancer tests and implications for cancer prevention, interspecies extrapolation and regulatory policy. The Carcinogenic Potency Database (CPDB), published as a CRC handbook, analyzes results of 5000 chronic, long-term cancer tests on 1300 chemicals. Dr. Gold has served on the Panel of Expert Reviewers for the National Toxicology Program, the Boards of the Harvard Center for Risk Analysis and the Annapolis Center, and was a member of the Harvard Risk Management Group. Lois@potency.Berkeley.edu (510)486-7080

Summary

1. The major causes of cancer are:

- a) **Smoking: About a third of U.S. cancer (90% of lung cancer);**
- b) **Dietary imbalances, e.g., lack of dietary fruits & vegetables: The quarter of the population eating the least fruits & vegetables has double the cancer rate for most types of cancer compared to the quarter eating the most;**
- c) **Chronic infections: mostly in developing countries;**
- d) **Hormonal factors: primarily influenced by life style.**

2. There is no epidemic of cancer, except for lung cancer due to smoking. Cancer mortality rates have declined 16% since 1950 (excluding lung cancer).

3. Regulatory policy that focuses on traces of synthetic chemicals is based on misconceptions about animal cancer tests. Recent research indicates that:

- a) **Rodent carcinogens are not rare. Half of all chemicals tested in standard high dose animal cancer tests, whether occurring naturally or produced synthetically, are "carcinogens";**
- b) **There are high-dose effects in rodent cancer tests that are not relevant to low-dose human exposures and which contribute to the high proportion of chemicals that test positive;**
- c) **The focus of regulatory policy is on synthetic chemicals, although 99.9% of the chemicals humans ingest are natural.**

*** Over 1000 chemicals have been described in coffee: 28 have been tested and 19 are rodent carcinogens.**

*** Plants in the human diet contain thousands of natural pesticides which protect them from insects and other predators: 63 have been tested and 35 are rodent carcinogens.**

4. There is no convincing evidence that synthetic chemical pollutants are important for human cancer. Regulations that try to eliminate minuscule levels of synthetic chemicals are enormously expensive: EPA has estimated that environmental regulations cost society \$140 billion/year. It has been estimated that the median toxic control program costs 146 times more per life year saved than the median medical intervention. Attempting to reduce tiny hypothetical risks also has costs, e.g., if reducing synthetic pesticides makes fruits and vegetables more expensive, thereby decreasing consumption, then cancer will be increased, particularly for the poor.

5. Prevention of cancer will come from knowledge obtained from biomedical research, education of the public, and from lifestyle changes by individuals. A re-examination of priorities in cancer prevention, both public and private, seems called for.

*Modified from testimony (3/6/97) for the U.S. Senate Hearing on Environmental Risk Factors for Cancer.

Various misconceptions about the relationship between environmental pollution and human disease, particularly cancer, drive regulatory policy. We highlight nine such misconceptions and briefly present the scientific evidence that undermines each.

Misconception #1: Cancer rates are soaring. Cancer death rates overall in the U.S. (excluding lung cancer due to smoking) have declined 16% since 1950 (1). The types of cancer deaths that have decreased since 1950 are primarily stomach, cervical, uterine, and colorectal. The types that have increased are primarily lung cancer (90% is due to smoking, as are 35 % of all cancer deaths in the U.S.), melanoma (probably due to sunburns), and non-Hodgkin's lymphoma. If lung cancer is included, mortality rates have increased over time, but recently have declined in men due to decreased smoking (1). The rise in incidence rates in older age groups for some cancers, e.g., prostate, can be explained by known factors such as improved screening. "The reason for not focusing on the reported incidence of cancer is that the scope and precision of diagnostic information, practices in screening and early detection, and criteria for reporting cancer have changed so much over time that trends in incidence are not reliable" (2). (See also (3) and (4)). Life expectancy has continued to rise since 1950.

Misconception #2: Environmental synthetic chemicals are an important cause of human cancer. Neither epidemiology nor toxicology supports the idea that synthetic industrial chemicals are important for human cancer (4-6). Epidemiological studies have identified the factors that are likely to have a major effect on lowering rates of cancer: reduction of smoking, improving diet (e.g., increased consumption of fruits and vegetables), hormonal factors and control of infections (6). Although some epidemiologic studies find an association between cancer and low levels of industrial pollutants, the associations are usually weak, the results are usually conflicting, and the studies do not correct for potentially large confounding factors like diet. Moreover, exposures to synthetic pollutants are tiny and rarely seem toxicologically plausible as a causal factor, particularly when compared to the background of natural chemicals that are rodent carcinogens (5). Even assuming that worst-case risk estimates for synthetic pollutants are true risks, the proportion of cancer that EPA could prevent by regulation would be tiny (7). Occupational exposures to some carcinogens cause cancer, though how much has been a controversial issue: a few percent seems a reasonable estimate (6), much of this from asbestos in smokers. Exposures to substances in the workplace can be high in comparison with other chemical exposures in food, air, or water. Past occupational exposures have sometimes been high and therefore comparatively little quantitative extrapolation may be required for risk assessment from high-dose rodent tests to high-dose occupational exposures. Since occupational cancer is concentrated among small groups exposed at high levels, there is an opportunity to control or eliminate risks once they are identified; however, current permitted workplace exposures are sometimes close to the carcinogenic dose in rodents (8).

Cancer is due, in part, to normal aging and increases exponentially with age in both rodents and humans (9). To the extent that the major external risk factors for cancer are diminished, cancer will occur at a later age, and the proportion of cancer caused by normal metabolic processes will increase. Aging and its degenerative diseases appear to be due in good part to oxidative damage to DNA and other macromolecules (9). By-products of normal metabolism -- superoxide, hydrogen peroxide, and hydroxyl radical -- are the same oxidative mutagens produced by radiation. Mitochondria from old animals leak oxidants (10); old rats have about 150,000 oxidative DNA lesions per cell (9, 11). DNA is oxidized in normal metabolism because antioxidant defenses, though numerous, are not perfect. Antioxidant defenses against oxidative damage include Vitamins C and E and perhaps carotenoids (12), most of which come from dietary fruits and vegetables.

Smoking contributes to about 35% of U.S. cancer, about one-quarter of heart disease, and about 400,000 premature deaths per year in the United States (6, 13). Tobacco is a known cause of cancer of the lung, bladder, mouth, pharynx, pancreas, stomach, larynx, esophagus and possibly colon. Tobacco causes even more deaths by diseases other than cancer. Smoke contains a wide variety of mutagens and rodent carcinogens. Smoking is also a severe oxidative stress and causes inflammation in the lung. The oxidants in cigarette smoke--mainly nitrogen oxides--deplete the body's antioxidants. Thus, smokers must ingest two to three times more Vitamin C than non-smokers to achieve the same level in blood, but they rarely do. Inadequate concentration of Vitamin C in plasma is more common among the poor and smokers. Men with inadequate diets or who smoke may damage both their somatic DNA and the DNA of their sperm. When the level of dietary Vitamin C is insufficient to keep seminal fluid Vitamin C at an adequate level, the oxidative lesions in sperm DNA are increased 250% (14-16). Male smokers, compared to non-smokers, have more oxidative lesions in sperm DNA (16) and more chromosomal abnormalities in sperm (17). Smoking by fathers, therefore, may plausibly increase the risk of birth defects and childhood cancer in offspring (14, 15, 18). A new epidemiological study suggests that the rate of childhood cancers is increased in offspring of male smokers, e.g., acute lymphocytic leukemia, lymphoma, and brain tumors, are increased 3-4 times (19).

We (6) estimate that unbalanced diets account for about one-third of cancer risk, in agreement with the earlier estimate of Doll and Peto (3). Low intake of fruits and vegetables is a major risk factor for cancer (See Misconception #3). There has been considerable interest in calories (and dietary fat) as a risk factor for cancer, in part because caloric restriction markedly lowers the cancer rate and increases life span in rodents (6, 20, 21).

Chronic inflammation from chronic infection results in release of oxidative mutagens from phagocytic cells and is a major contributor to cancer (6, 22). White cells and other phagocytic cells of the immune system combat bacteria, parasites, and virus-infected cells by destroying them with potent, mutagenic oxidizing agents. The oxidants protect humans from immediate death from infection, but they also cause oxidative damage to DNA, chronic cell killing with compensatory cell division, and mutation (23, 24) and thus contribute to the carcinogenic process. Antioxidants appear to inhibit some of the pathology of chronic inflammation. Chronic infections cause about 21% of new cancer cases in developing countries and 9% in developed countries (25).

Endogenous reproductive hormones play a large role in cancer, including breast, prostate, ovary and endometrium (26, 27), contributing to as much as 20% of all cancer. Many lifestyle factors such as reproductive history, lack of exercise, obesity, and alcohol influence hormone levels and therefore increase risk (6, 26-28).

Other causal factors in human cancer are excessive alcohol consumption, excessive sun exposure, and viruses. Genetic factors also play a significant role and interact with lifestyle and other risk factors. Biomedical research is uncovering important genetic variation in humans.

Misconception #3: Reducing pesticide residues is an effective way to prevent diet-related cancer. On the contrary, fruits and vegetables are of major importance for reducing cancer; if they become more expensive because of reduced use of synthetic pesticides, then cancer is likely to increase. People with low incomes eat fewer fruits and vegetables and spend a higher percentage of their income on food.

Dietary fruits and vegetables and cancer prevention. High consumption of fruits and vegetables is associated with a lowered risk of degenerative diseases including cancer, cardiovascular disease, cataracts, and brain dysfunction (6, 9). Over 200 studies in the epidemiological literature have been reviewed that show, with great consistency, an association between low consumption of fruits and vegetables and cancer incidence (29-31) (Table 1). The quarter of the population with the lowest dietary intake of fruits and vegetables compared to the quarter with the highest

intake has roughly twice the cancer rate for most types of cancer (lung, larynx, oral cavity, esophagus, stomach, colorectal, bladder, pancreas, cervix, and ovary). Only 22% of Americans met the intake recommended by the NCI and the National Research Council (32, 33): 5 servings of fruits and vegetables per day. Publicity about hundreds of minor hypothetical risks can cause loss of perspective on what is important: half the public does not know that fruit and vegetable consumption is a major protection against cancer (34).

Some micronutrients in fruits and vegetables are anticarcinogens. Antioxidants in fruits and vegetables may account for some of their beneficial effect, as discussed in Misconception #2. However, the effects of dietary antioxidants are difficult to disentangle by epidemiological studies from other important vitamins and ingredients in fruits and vegetables (30, 31, 35).

Folate deficiency, one of the most common vitamin deficiencies, causes extensive chromosome breaks in human genes (36). Approximately 10% of the US population (37) has a lower folate level than that at which chromosome breaks occur (36). In two small studies of low income (mainly African-American) elderly (38) and adolescents (39) nearly half had folate levels that low. The mechanism is deficient methylation of uracil to thymine, and subsequent incorporation of uracil into human DNA (4 million/cell) (36). During repair of uracil in DNA, transient nicks are formed; two opposing nicks causes a chromosome break. Both high DNA uracil levels and chromosome breaks in humans are reversed by folate administration (36). Chromosome breaks could contribute to the increased risk of cancer and cognitive defects associated with folate deficiency in humans (36). Folate deficiency also damages human sperm (40), causes neural tube defects in the fetus, and about 10% of the risk of U.S. heart disease (41).

Other micronutrients whose main dietary sources are other than fruits and vegetables, also are likely to play a significant role in the prevention and repair of DNA damage, and thus are important to the maintenance of long term health. Deficiency of Vitamin B₁₂ causes a functional folate deficiency, accumulation of homocysteine (a risk factor for heart disease) (42), and misincorporation of uracil into DNA (43). Strict vegetarians are at increased risk of developing a Vitamin B₁₂ deficiency (42). Niacin contributes to the repair of DNA strand breaks by maintaining nicotinamide adenine dinucleotide levels for the poly ADP-ribose protective response to DNA damage (44). As a result, dietary insufficiencies of niacin (15-20% of several populations are deficient (45)), folate, and antioxidants may act synergistically to adversely affect DNA synthesis and repair. Diets deficient in fruits and vegetables are commonly low in folate, antioxidants, (e.g., Vitamin C) and many other micronutrients, and result in DNA damage and higher cancer rates (6, 29, 46).

Optimizing micronutrient intake can have a major impact on health at low cost. Increasing research in this area and efforts to increase micronutrient intake and more balanced diets, should be high priorities for public policy.

Misconception #4: Human exposures to carcinogens and other potential hazards are primarily to synthetic chemicals. On the contrary, 99.9% of the chemicals humans ingest are natural. The amounts of synthetic pesticide residues in plant foods are insignificant compared to the amount of natural pesticides produced by plants themselves (47-49). Of all dietary pesticides that humans eat, 99.99% are natural: they are chemicals produced by plants to defend themselves against fungi, insects, and other animal predators (47, 48). Each plant produces a different array of such chemicals. On average Americans ingest roughly 5,000 to 10,000 different natural pesticides and their breakdown products. Americans eat about 1,500 mg of natural pesticides per person per day, which is about 10,000 times more than they consume of synthetic pesticide residues.

Even though only a small proportion of natural pesticides has been tested for carcinogenicity, half of those tested (35/63) are rodent carcinogens, and naturally occurring pesticides that are rodent carcinogens are ubiquitous in fruits, vegetables, herbs, and spices (49) (Table 2).

Cooking foods produces about 2,000 mg per person per day of burnt material that contains many rodent carcinogens and many mutagens. By contrast, the residues of 200 synthetic chemicals measured by FDA, including the synthetic pesticides thought to be of greatest importance, average only about 0.09 mg per person per day (47, 49). In a single cup of coffee the known natural chemicals that are rodent carcinogens, are about equal in weight to an entire year's worth of synthetic pesticide residues that are rodent carcinogens, even though only 3% of the natural chemicals in roasted coffee have been adequately tested for carcinogenicity (5) (Table 3). This does not mean that coffee or natural pesticides are dangerous, but rather that assumptions about high dose animal cancer tests for assessing human risk at low doses need reexamination. No diet can be free of natural chemicals that are rodent carcinogens (49).

Misconception #5: Cancer risks to humans can be assessed by standard high-dose animal cancer tests. Approximately half of all chemicals -- whether natural or synthetic -- that have been tested in standard animal cancer tests are rodent carcinogens (50, 51) (Table 4). What are the explanations for the high positivity rate? In standard cancer tests rodents are given chronic, near-toxic doses, the maximum tolerated dose (MTD). Evidence is accumulating that cell division caused by the high dose itself, rather than the chemical per se, is increasing the positivity rate. High doses can cause chronic wounding of tissues, cell death, and consequent chronic cell division of neighboring cells, which is a risk factor for cancer (50). Each time a cell divides the probability increases that a mutation will occur, thereby increasing the risk for cancer. At the low levels to which humans are usually exposed, such increased cell division does not occur. In addition, tissues injured by high doses of chemicals (e.g., phenobarbital, carbon tetrachloride, TPA) have an inflammatory immune response involving activation of recruited and resident macrophages in response to necrosis (52-58). Activated macrophages release mutagenic oxidants (including peroxynitrite, hypochlorite, and H_2O_2). Therefore, the very low levels of chemicals to which humans are exposed through water pollution or synthetic pesticide residues may pose no or minimal cancer risks.

We have discussed (59) the argument that the high positivity rate is due to selecting more suspicious chemicals to test, which is a likely bias since cancer testing is both expensive and time-consuming, and it is prudent to test suspicious compounds. One argument against selection bias is the high positivity rate for drugs (Table 4) because drug development tends to select chemicals which are not mutagens or expected carcinogens. A second argument against selection bias is that knowledge to predict carcinogenicity in rodent tests is highly imperfect, even now after decades of testing results have become available on which to base prediction. For example, a prospective prediction exercise was conducted by several experts in 1990 in advance of the 2-year NTP bioassays. There was wide disagreement among them on which chemicals would be carcinogenic when tested, and accuracy varied, thus indicating that predictive knowledge is highly uncertain (60). Moreover, if the main basis for selection was suspicion rather than human exposure, then one should select mutagens (79% are positive compared to 49% of non-mutagens), yet 55% of the chemicals tested are non-mutagens (59).

It seems likely that a high proportion of all chemicals, whether synthetic or natural, might be "carcinogens" if run through the standard rodent bioassay at the MTD, but this will be primarily due to the effects of high doses for the non-mutagens, and for the mutagens, a synergistic effect between cell division at high doses and DNA damage (61-63). Without additional data on mechanism of carcinogenesis for each chemical, the interpretation of a positive result in a rodent bioassay is highly uncertain. The carcinogenic effects may be limited to the high dose tested.

In regulatory policy, the "virtually safe dose" (VSD), corresponding to a maximum, hypothetical cancer risk of one in a million, is estimated from bioassay results using a linear model. To the extent that carcinogenicity in rodent bioassays is due to the effects of high doses for the non-mutagens, and a synergistic effect of cell division at high doses with DNA damage for the mutagens, then this model is

inappropriate. Moreover, as currently calculated, the VSD can be known without ever conducting a bioassay: for 96% of the NCI/NTP rodent carcinogens, the VSD is within a factor of 10 of the ratio MTD/740,000 (64). This is about as precise as the estimate obtained from conducting near-replicate cancer tests of the same chemical (64).

Misconception #6: Synthetic chemicals pose greater carcinogenic hazards than natural chemicals. Gaining a broad perspective about the vast number of chemicals to which humans are exposed can be helpful when setting research and regulatory priorities (5, 48, 49, 65). Rodent bioassays provide little information about mechanisms of carcinogenesis and low-dose risk. The assumption that synthetic chemicals are hazardous has led to a bias in testing, such that synthetic chemicals account for 77% (432/559) of the chemicals tested chronically in both rats and mice (Table 4). The natural world of chemicals has never been tested systematically. One reasonable strategy is to use a rough index to *compare* and *rank* possible carcinogenic hazards from a wide variety of chemical exposures at levels that humans typically receive, and then to focus on those that rank highest (5, 51). Ranking is a critical first step that can help to set priorities for selecting chemicals for chronic bioassay or mechanistic studies, for epidemiological research, and for regulatory policy. Although one cannot say whether the ranked chemical exposures are likely to be of major or minor importance in human cancer, it is not prudent to focus attention on the possible hazards at the bottom of a ranking if, using the same methodology to identify hazard, there are numerous common human exposures with much greater possible hazards. Our analyses are based on the HERP index (Human Exposure/Rodent Potency), which indicates what percentage of the rodent carcinogenic potency (TD₅₀ in mg/kg/day) a human receives from a given daily lifetime exposure (mg/kg/day) (66) (Table 5). A ranking based on standard regulatory risk assessment would be similar.

Overall, our analyses have shown that HERP values for some historically high exposures in the workplace and some pharmaceuticals rank high, and that there is an enormous background of naturally occurring rodent carcinogens in typical portions of common foods that cast doubt on the relative importance of low-dose exposures to residues of synthetic chemicals such as pesticides (5, 8, 51). A committee of the National Research Council/National Academy of Sciences recently reached similar conclusions about natural vs. synthetic chemicals in the diet, and called for further research on natural chemicals (67).

The possible carcinogenic hazards from synthetic pesticides (at average exposures) are minimal compared to the background of nature's pesticides, though neither may be a hazard at the low doses consumed (Table 5). Table 5 also indicates that many ordinary foods would not pass the regulatory criteria used for synthetic chemicals. For many natural chemicals the HERP values are in the top half of the table, even though natural chemicals are markedly underrepresented because so few have been tested in rodent bioassays. Caution is necessary in drawing conclusions from the occurrence in the diet of natural chemicals that are rodent carcinogens. It is not argued here that these dietary exposures are necessarily of much relevance to human cancer. Our results call for a re-evaluation of the utility of animal cancer tests in protecting the public against minor hypothetical risks.

Misconception #7: The toxicology of synthetic chemicals is different from that of natural chemicals. It is often assumed that because natural chemicals are part of human evolutionary history, whereas synthetic chemicals are recent, the mechanisms that have evolved in animals to cope with the toxicity of natural chemicals will fail to protect against synthetic chemicals. This assumption is flawed for several reasons (48, 50).

a) Humans have many natural defenses that buffer against normal exposures to toxins (48), and these are usually general, rather than tailored for each specific chemical. Thus they work against both natural and synthetic chemicals. Examples of general defenses include the continuous shedding of cells exposed to toxins -- the

surface layers of the mouth, esophagus, stomach, intestine, colon, skin, and lungs are discarded every few days; DNA repair enzymes, which repair DNA that was damaged from many different sources; and detoxification enzymes of the liver and other organs which generally target classes of toxins rather than individual toxins. That defenses are usually general, rather than specific for each chemical, makes good evolutionary sense. The reason that predators of plants evolved general defenses is presumably to be prepared to counter a diverse and ever-changing array of plant toxins in an evolving world; if a herbivore had defenses against only a set of specific toxins, it would be at a great disadvantage in obtaining new food when favored foods became scarce or evolved new toxins.

b) Various natural toxins, which have been present throughout vertebrate evolutionary history, nevertheless cause cancer in vertebrates (48, 51). Mold toxins, such as aflatoxin, have been shown to cause cancer in rodents and other species including humans (Table 4). Many of the common elements are carcinogenic to humans at high doses (e.g., salts of cadmium, beryllium, nickel, chromium, and arsenic) despite their presence throughout evolution. Furthermore, epidemiological studies from various parts of the world show that certain natural chemicals in food may be carcinogenic risks to humans; for example, the chewing of betel nuts with tobacco has been correlated with oral cancer.

c) Humans have not had time to evolve a "toxic harmony" with all of their dietary plants. The human diet has changed markedly in the last few thousand years. Indeed, very few of the plants that humans eat today (e.g., coffee, cocoa, tea, potatoes, tomatoes, corn, avocados, mangoes, olives, and kiwi fruit), would have been present in a hunter-gatherer's diet. Natural selection works far too slowly for humans to have evolved specific resistance to the food toxins in these newly introduced plants.

d) DDT is often viewed as the typically dangerous synthetic pesticide because it concentrates in the tissues and persists for years, being slowly released into the bloodstream. DDT, the first synthetic pesticide, eradicated malaria from many parts of the world, including the U.S. It was effective against many vectors of disease such as mosquitoes, tsetse flies, lice, ticks, and fleas. DDT was also lethal to many crop pests, and significantly increased the supply and lowered the cost of food, making fresh nutritious foods more accessible to poor people. It was also remarkably non-toxic to humans. A 1970 National Academy of Sciences report concluded: "In little more than two decades DDT has prevented 500 million deaths due to malaria, that would otherwise have been inevitable (68)." There is no convincing epidemiological evidence, nor is there much toxicological plausibility, that the levels normally found in the environment are likely to be a significant contributor to cancer. DDT was unusual with respect to bioconcentration, and because of its chlorine substituents it takes longer to degrade in nature than most chemicals; however, these are properties of relatively few synthetic chemicals. In addition, many thousands of chlorinated chemicals are produced in nature (69) and natural pesticides also can bioconcentrate if they are fat soluble. Potatoes, for example, naturally contain the fat soluble neurotoxins solanine and chaconine (49), which can be detected in the bloodstream of all potato eaters. High levels of these potato neurotoxins have been shown to cause birth defects in rodents (48).

e) Since no plot of land is immune to attack by insects, plants need chemical defenses -- either natural or synthetic -- in order to survive pest attack. Thus, there is a trade-off between naturally occurring pesticides and synthetic pesticides. One consequence of disproportionate concern about synthetic pesticide residues is that some plant breeders develop plants to be more insect-resistant by making them higher in natural toxins. A recent case illustrates the potential hazards of this approach to pest control: When a major grower introduced a new variety of highly insect-resistant celery into commerce, people who handled the celery developed rashes when they were subsequently exposed to sunlight. Some detective work found that the pest-resistant celery contained 6,200 parts per billion (ppb) of carcinogenic (and mutagenic) psoralens instead of the 800 ppb present in common celery (49).

Misconception #8: Pesticides and other synthetic chemicals are disrupting hormones. Synthetic hormone mimics have become an environmental issue. Hormonal factors are important in cancer (Misconception #2). A recent book (70), states that traces of synthetic chemicals, such as pesticides with weak hormonal activity, may contribute to cancer and reduce sperm counts. The book ignores the fact that our normal diet contains natural chemicals that have estrogenic activity millions of times higher than that due to the traces of synthetic estrogenic chemicals (71, 72) and that lifestyle factors can markedly change the levels of endogenous hormones (Misconception #2). The low levels of exposure to residues of industrial chemicals in humans are toxicologically implausible as a significant cause of cancer or of reproductive abnormalities, especially when compared to the natural background (71-74). In addition, it has not been shown convincingly that sperm counts are declining (75), and even if they were, there are many more likely causes, such as smoking and diet (Misconception # 2).

Misconception #9: Regulation of low, hypothetical risks is effective in advancing public health. Since there is no risk-free world and resources are limited, society must set priorities based on cost effectiveness in order to save the most lives (76, 77). The EPA projected in 1991 that the cost to society of environmental regulations in 1997 would be about \$140 billion per year (about 2.6% of GNP) (78). Most of this cost is to the private sector. Several economic analyses have concluded that current expenditures are not cost effective, i.e. resources are not being utilized so as to save the most lives per dollar. One estimate is that the U.S. could prevent 60,000 deaths per year by redirecting the same dollar resources to more cost effective programs (79). For example, the median toxin control program costs 146 times more per life-year saved than the median medical intervention (79). This difference is likely to be greater, because cancer risk estimates for toxin control programs are worst-case, hypothetical estimates, and the true risks at low dose are often likely to be zero (5, 6, 51) (Misconception #5). Some economists have argued that costly regulations intended to save lives may actually lead to increased deaths (80), in part because they divert resources from important health risks and in part because higher incomes are associated with lower mortality (81-83). Rules on air and water pollution are necessary (e.g., it was a public health advance to phase lead out of gasoline) and clearly, cancer prevention is not the only reason for regulations. However, worst-case assumptions in risk assessment represent a policy decision, not a scientific one, and they confuse attempts to allocate money effectively for risk abatement.

Regulatory efforts to reduce low-level human exposures to synthetic chemicals because they are rodent carcinogens, are expensive because they aim to eliminate minuscule concentrations that now can be measured with improved techniques. These efforts are distractions from the major task of improving public health through increasing scientific understanding about how to prevent cancer (e.g., the role of diet), increasing public understanding of how lifestyle influences health, and improving our ability to help individuals alter lifestyle.

Acknowledgments. This work was supported by the National Cancer Institute Outstanding Investigator Grant CA39910 to B.N.A., the Director, Office of Energy Research, Office of Health and Environmental Research of the U.S. Department of Energy under Contract DE-AC03-76SF00098 to L.S.G., and the National Institute of Environmental Health Sciences Center Grant ESO1896.

1. Ries, L.A.G., *et al.*, *SEER Cancer Statistics Review, 1973-1994*. 1997, Bethesda, MD: National Cancer Institute.
2. Bailar, I., J C and H.L. Gornik, *Cancer undefeated*. N. Engl. J. Med., 1997. **336**: p. 1569-1574.
3. Doll, R. and R. Peto, *The causes of cancer. Quantitative estimates of avoidable risks of cancer in the United States today*. J. Natl. Cancer Inst., 1981. **66**: p. 1191-1308.
4. Devesa, S.S., *et al.*, *Recent cancer trends in the United States*. J. Natl. Cancer Inst., 1995. **87**: p. 175-182.
5. Gold, L.S., *et al.*, *Rodent carcinogens: Setting priorities*. Science, 1992. **258**: p. 261-265.
6. Ames, B.N., L.S. Gold, and W.C. Willett, *The causes and prevention of cancer*. Proc. Natl. Acad. Sci. USA, 1995. **92**: p. 5258-5265.
7. Gough, M., *How much cancer can EPA regulate anyway?* Risk Anal., 1990. **10**: p. 1-6.
8. Gold, L.S., G.B. Garfinkel, and T.H. Slone, *Setting priorities among possible carcinogenic hazards in the workplace, in Chemical Risk Assessment and Occupational Health, Current Applications, Limitations, and Future Prospects*, C.M. Smith, D.C. Christiani, and K.T. Kelsey, Editors. 1994, Greenwood Publishing Group: Westport, CT. p. 91-103.
9. Ames, B.N., M.K. Shigenaga, and T.M. Hagen, *Oxidants, antioxidants, and the degenerative diseases of aging*. Proc. Natl. Acad. Sci. USA, 1993. **90**: p. 7915-7922.
10. Hagen, T.M., *et al.*, *Mitochondrial decay in hepatocytes from old rats: Membrane potential declines, heterogeneity and oxidants increase*. Proc. Natl. Acad. Sci. USA, 1997. **94**: p. 3064-3069.
11. Beckman, K.B. and B.N. Ames, *Oxidative decay of DNA*. J. Biol. Chem., 1997. **272**: p. 19633-19636.
12. Rice-Evans, C., *et al.*, *Why do we expect carotenoids to be antioxidants in vivo?* Free Rad. Res., 1997. **26**: p. 381-398.
13. Peto, R., *et al.*, *Mortality from Smoking in Developed Countries 1950-2000*. 1994, Oxford, England: Oxford University Press.
14. Fraga, C.G., *et al.*, *Ascorbic acid protects against endogenous oxidative damage in human sperm*. Proc. Natl. Acad. Sci. USA, 1991. **88**: p. 11003-11006.
15. Ames, B.N., *et al.*, *Antioxidant prevention of birth defects and cancer, in Male-Mediated Developmental Toxicity*, D.R. Mattison and A. Olshan, Editors. 1994, Plenum Publishing Corporation: New York. p. 243-259.
16. Fraga, C.G., *et al.*, *Smoking and low antioxidant levels increase oxidative damage to sperm DNA*. Mutat. Res., 1996. **351**: p. 199-203.
17. Wyrobek, A.J., *et al.*, *Smokers produce more aneuploid sperm than non-smokers*. Am. J. Hum. Genet., 1995. **57**: p. 737.
18. Woodall, A.A. and B.N. Ames, *Diet and oxidative damage to DNA: The importance of ascorbate as an antioxidant, in Vitamin C in Health and Disease*, L. Packer, Editor. 1997, Marcel Dekker, Inc.: New York. p. 193-203.
19. Ji, B.-T., *et al.*, *Paternal cigarette smoking and the risk of childhood cancer among offspring of nonsmoking mothers*. J. Natl. Cancer Inst., 1997. **89**: p. 238-244.
20. Hart, R., *et al.*, *Caloric restriction and toxicity*. Fundam. Appl. Toxicol., 1995. **25**: p. 184-195.
21. Turturro, A., *et al.*, *Rationale for the use of dietary control in toxicity studies--B6C3F1 mouse*. Toxicol. Pathol., 1996. **24**: p. 769-775.
22. Christen, S., *et al.*, *Chronic infection and inflammation lead to mutation and cancer, in Microbes and Malignancy: Infection as a Cause of Cancer*, J. Parsonnet and S. Horning, Editors. 1997, Oxford University Press: Oxford. p. in press.
23. Shacter, E., *et al.*, *Activated neutrophils induce prolonged DNA damage in neighboring cells [published erratum appears in Carcinogenesis 1989 Mar;10(3):628]*. Carcinogenesis, 1988. **9**(12): p. 2297-2304.
24. Yamashina, K., B.E. Miller, and G.H. Heppner, *Macrophage-mediated induction of drug-resistant variants in a mouse mammary tumor cell line*. Cancer Res., 1986. **46**: p. 2396-2401.
25. Pisani, P., *et al.*, *Cancer and infection: Estimates of the attributable fraction in 1990*. Cancer Epidemiol. Biomarkers & Prev, 1997. **6**: p. 387-400.
26. Henderson, B.E., R.K. Ross, and M.C. Pike, *Toward the primary prevention of cancer*. Science, 1991. **254**: p. 1131-1138.
27. Feigelson, H.S. and B.E. Henderson, *Estrogens and breast cancer*. Carcinogenesis, 1996. **17**: p. 2279-2284.
28. Hunter, D.J. and W.C. Willett, *Diet, body size, and breast cancer*. Epidemiol. Rev., 1993. **15**(1): p. 110-132.
29. Block, G., B. Patterson, and A. Subar, *Fruit, vegetables and cancer prevention: A review of the epidemiologic evidence*. Nutr. and Cancer, 1992. **18**: p. 1-29.
30. Steinmetz, K.A. and J.D. Potter, *Vegetables, fruit, and cancer prevention: A review*. J. Am. Diet Assoc., 1996. **96**: p. 1027-1039.
31. Hill, M.J., A. Giacosa, and C.P.J. Caygill, *Epidemiology of Diet and Cancer*. Ellis Horwood Series in Food Science and Technology, ed. D.H. Watson and M.J. Lewis. 1994, West Sussex, England: Ellis Horwood Limited.
32. Block, G., *Epidemiologic evidence regarding vitamin C and cancer*. Am. J. Clin. Nutr., 1991. **54**: p. 1310S-1314S.
33. Patterson, B.H., *et al.*, *Fruit and vegetables in the American diet: Data from the NHANES II survey*. Am. J. Public Health, 1990. **80**: p. 1443-1449.
34. A National Cancer Institute Graphic, *Why eat five?* J. Natl. Cancer Inst., 1996. **88**: p. 1314.
35. Block, G., *The data support a role for antioxidants in reducing cancer risk*. Nutr. Reviews, 1992. **50**: p. 207-213.
36. Blount, B.C., *et al.*, *Folate deficiency causes uracil misincorporation into human DNA and chromosome breakage: Implications for cancer and neuronal damage*. Proc Natl Acad Sci USA, 1997. **94**: p. 3290-3295.
37. Senti, F.R. and S.M. Pilch, *Analysis of folate data from the second National Health and Nutrition Examination Survey (NHANES II)*. J. Nutr., 1985. **115**(11): p. 1398-402.
38. Bailey, L.B., *et al.*, *Folacin and iron status and hematological findings in predominately black elderly persons from urban low-income households*. Am. J. Clin. Nutr., 1979. **32**(11): p. 2346-2353.
39. Bailey, L.B., *et al.*, *Folacin and iron status and hematological findings in black and Spanish-American adolescents from urban low-income households*. Am. J. Clin. Nutr., 1982. **35**: p. 1023-1032.

40. Wallock, L., *et al.* Nutritional status and positive relation of plasma folate to fertility indices in nonsmoking men. in *FASEB Annual Meeting, Experimental Biology* 97. 1997. New Orleans, LA: The FASEB J.
41. Boushey, C.J., *et al.*, A quantitative assessment of plasma homocysteine as a risk factor for vascular disease. *Probable benefits of increasing folic acid intakes.* *Jama*, 1995. **274**(13): p. 1049-57.
42. Herbert, V., *Vitamin B-12*, in *Present Knowledge in Nutrition*, E.E. Ziegler and L.J. Filer, Editors. 1996, ILSI Press: Washington, D.C. p. 191-205.
43. Wickramasinghe, S.N. and S. Fida, *Bone marrow cells from vitamin B12- and folate-deficient patients misincorporate uracil into DNA.* *Blood*, 1994. **83**(6): p. 1656-61.
44. Zhang, J.Z., S.M. Henning, and M.E. Swendseid, *Poly(ADP-ribose) polymerase activity and DNA strand breaks are affected in tissues of niacin-deficient rats.* *J. Nutr.*, 1993. **123**: p. 1349-55.
45. Jacobson, E.L., *Niacin deficiency and cancer in women.* *J. Am. Coll. Nutr.*, 1993. **12**(4): p. 412-6.
46. Subar, A.F., G. Block, and L.D. James, *Folate intake and food sources in the US population.* *Am. J. Clin. Nutr.*, 1989. **50**(3): p. 508-16.
47. Ames, B.N., M. Profet, and L.S. Gold, *Dietary pesticides (99.99% all natural).* *Proc. Natl. Acad. Sci. USA*, 1990. **87**: p. 7777-7781.
48. Ames, B.N., M. Profet, and L.S. Gold, *Nature's chemicals and synthetic chemicals: Comparative toxicology.* *Proc. Natl. Acad. Sci. USA*, 1990. **87**: p. 7782-7786.
49. Gold, L.S., T.H. Slone, and B.N. Ames, *Prioritization of possible carcinogenic hazards in food*, in *Food Chemical Risk Analysis*, D. Tennant, Editor. 1997, Chapman & Hall Ltd: London. p. in press.
50. Ames, B.N., L.S. Gold, and M.K. Shigenaga, *Cancer prevention, rodent high-dose cancer tests and risk assessment.* *Risk Anal.*, 1996. **16**: p. 613-617.
51. Gold, L.S., T.H. Slone, and B.N. Ames, *Overview and update analyses of the carcinogenic potency database*, in *Handbook of Carcinogenic Potency and Genotoxicity Databases*, L.S. Gold and E. Zeiger, Editors. 1997, CRC Press: Boca Raton, FL. p. 661-685.
52. Laskin, D.L. and K.J. Pendino, *Macrophages and inflammatory mediators in tissue injury.* *Annu Rev Pharmacol Toxicol*, 1995. **35**(655): p. 655-677.
53. Wei, H. and K. Frenkel, *Relationship of oxidative events and DNA oxidation in SENCAR mice to in vivo promoting activity of phorbol ester-type tumor promoters.* *Carcinogenesis*, 1993. **14**(6): p. 1195-1201.
54. Wei, L., H. Wei, and K. Frenkel, *Sensitivity to tumor promotion of SENCAR and C57BL/6J mice correlates with oxidative events and DNA damage.* *Carcinogenesis*, 1993. **14**(5): p. 841-847.
55. Laskin, D.L., *et al.*, *Activation of liver macrophages following phenobarbital treatment of rats.* *Hepatology*, 1988. **8**(5): p. 1051-1055.
56. Czaja, M.J., *et al.*, *Lipopolysaccharide-neutralizing antibody reduces hepatocyte injury from acute hepatotoxin administration.* *Hepatology*, 1994. **19**(5): p. 1282-1289.
57. Adachi, Y., *et al.*, *Antibiotics prevent liver injury in rats following long-term exposure to ethanol.* *Gastroenterology*, 1995. **108**(1): p. 218-224.
58. Gunawardhana, L., S.A. Mobley, and I.G. Sipes, *Modulation of 1,2-dichlorobenzene hepatotoxicity in the Fischer-344 rat by a scavenger of superoxide anions and an inhibitor of Kupffer cells.* *Toxicol Appl Pharmacol*, 1993. **119**(2): p. 205-213.
59. Gold, L.S., T.H. Slone, and B.N. Ames, *What do animal cancer tests tell us about human cancer risk?: Overview of analyses of the Carcinogenic Potency Database.* *Drug Metab. Rev.*, 1997: p. in press.
60. Omenn, G.S., S. Stuebbe, and L.B. Lave, *Predictions of rodent carcinogenicity testing results: interpretation in light of the Lave-Omenn value-of-information model.* *Mol. Carcinog.*, 1995. **14**: p. 37-45.
61. Butterworth, B., R. Conolly, and K. Morgan, *A strategy for establishing mode of action of chemical carcinogens as a guide for approaches to risk assessment.* *Cancer Lett.*, 1995. **93**: p. 129-146.
62. Ames, B.N., M.K. Shigenaga, and L.S. Gold, *DNA lesions, inducible DNA repair, and cell division: Three key factors in mutagenesis and carcinogenesis.* *Environ Health Perspect*, 1993. **101**(Suppl 5): p. 35-44.
63. Ames, B.N. and L.S. Gold, *Chemical carcinogenesis: Too many rodent carcinogens.* *Proc. Natl. Acad. Sci. USA*, 1990. **87**: p. 7772-7776.
64. Gaylor, D.W. and L.S. Gold, *Quick estimate of the regulatory virtually safe dose based on the maximum tolerated dose for rodent bioassays.* *Regul. Toxicol. Pharmacol.*, 1995. **22**: p. 57-63.
65. Ames, B.N., R. Magaw, and L.S. Gold, *Ranking possible carcinogenic hazards.* *Science*, 1987. **236**: p. 271-280.
66. Gold, L.S. and E. Zeiger, *Handbook of Carcinogenic Potency and Genotoxicity Databases.* 1997, Boca Raton, FL: CRC Press.
67. National Research Council, *Carcinogens and Anticarcinogens in the Human Diet: A Comparison of Naturally Occurring and Synthetic Substances*, . 1996, National Academy Press: Washington, D.C.
68. National Academy of Sciences, U.S.A., *The Life Sciences: Recent Progress and Application to Human Affairs, the World of Biological Research, Requirement for the Future*, . 1970, Committee on Research in the Life Sciences: Washington, D.C. p. 526 p.
69. Gribble, G.W., *The diversity of natural organochlorines in living organisms.* *Pure & Appl. Chem.*, 1996. **68**: p. 1699-1712.
70. Colburn, T., D. Dumanoski, and J.P. Myers, *Our Stolen Future: Are we Threatening our Fertility, Intelligence, and Survival?: A Scientific Detective Story.* 1996, New York: Dutton.
71. Safe, S.H., *Dietary and environmental estrogens and antiestrogens and their possible role in human disease.* *Environ. Sci. Pollution Res.*, 1994. **1**(29-33): p. 29-33.
72. Safe, S.H., *Environmental and dietary estrogens and human health: Is there a problem?* *Env. Health Persp.*, 1995. **103**: p. 346-351.
73. Safe, S.H., *Is there an association between exposure to environmental estrogens and breast cancer?* *Environ. Hlth. Persp.*, 1997. **105**: p. (Suppl 3): 675-678.

74. Reinli, K. and G. Block, *Phytoestrogen content of foods -- A compendium of literature values*. Nutr. Cancer, 1996. **26**(123-148): p. 1996.
75. Kolata, G., *Measuring men up, sperm by sperm*, in *The New York Times*,. 1996. p. E4(N), E4(L), (col.1).
76. Hahn, R.W., ed. *Risks, Costs, and Lives Saved: Getting Better Results from Regulation*. . 1996, Oxford University Press and AEI Press: New York and Washington, DC.
77. Graham, J. and J. Wiener, eds. *Risk versus Risk: Tradeoffs in Protecting Health and the Environment*. . 1995, Harvard University Press: Cambridge, Massachusetts.
78. U.S. Environmental Protection Agency, *Environmental Investments: The Cost of a Clean Environment*, . 1991, Office of the Administrator: Washington, D. C.
79. Tengs, T.O., *et al.*, *Five-hundred life-saving interventions and their cost-effectiveness*. Risk Anal., 1995. **15**(3): p. 369-389.
80. Keeney, R.L., *Mortality risks induced by economic expenditures*. Risk Anal., 1990. **10**: p. 147-159.
81. Wildavsky, A., *Searching for Safety*. 1988, New Brunswick, N.J.: Transaction Press.
82. Wildavsky, A.B., *But is it True?: A Citizen's Guide to Environmental Health and Safety*. 1995, Cambridge, MA: Harvard University Press.
83. Viscusi, W.K., *Fatal Trade-offs*. 1992, Oxford, England: Oxford University Press.
84. Innes, J.R.M., *et al.*, *Bioassay of pesticides and industrial chemicals for tumorigenicity in mice: A preliminary note*. J. Natl. Cancer Inst., 1969. **42**: p. 1101-1114.
85. Davies, T.S. and A. Monro, *Marketed Human Pharmaceuticals Reported to be Tumorigenic in Rodents*. J. Am. Coll. Toxicol., 1995. **14**(2): p. 90-107.
86. Contrera, J., A. Jacobs, and J. DeGeorge, *Carcinogenicity testing and the evaluation of regulatory requirements for pharmaceuticals*. Regul. Toxicol. Pharmacol., 1997. **25**: p. 130-145.

Table 1. Review of epidemiological studies on cancer showing protection by consumption of fruits and vegetables (29).

Cancer site	Fraction of studies showing significant cancer protection	Relative risk (median) Low vs. High Quartile) of consumption
Epithelial		
Lung	24/25	2.2
Oral	9/9	2.0
Larynx	4/4	2.3
Esophagus	15/16	2.0
Stomach	17/19	2.5
Pancreas	9/11	2.8
Cervix	7/8	2.0
Bladder	3/5	2.1
Colorectal	20/35	1.9
Miscellaneous	6/8	---
Hormone-dependent		
Breast	8/14	1.3
Ovary/endometrium	3/4	1.8
Prostate	4/14	1.3
Total	129/172	

Table 2. Carcinogenicity of natural plant pesticides tested in rodents (Fungal toxins are not included) (49).

Carcinogens: N=35	acetaldehyde methylformylhydrazone, allyl isothiocyanate, arecoline.HCl, benzaldehyde, benzyl acetate, caffeic acid, catechol, clivorine, coumarin, crotonaldehyde, cycasin and methylazoxymethanol acetate, 3,4-dihydrocoumarin, estragole, ethyl acrylate, <i>N</i> 2- γ -glutamyl- <i>p</i> -hydrazinobenzoic acid, hexanal methylformylhydrazine, <i>p</i> -hydrazinobenzoic acid.HCl, hydroquinone, 1-hydroxyanthraquinone, lasiocarpine, <i>d</i> -limonene, 8-methoxypsoralen, <i>N</i> -methyl- <i>N</i> -formylhydrazine, α -methylbenzyl alcohol, 3-methylbutanal methylformylhydrazone, methylhydrazine, monocrotaline, pentanal methylformylhydrazone, petasitenine, quercetin, reserpine, safrole, senkirkine, sesamol, symphytine
Noncarcinogens: N=28	atropine, benzyl alcohol, biphenyl, <i>d</i> -carvone, deserpidine, disodium glycyrrhizinate, emetine.2HCl, ephedrine sulphate, eucalyptol, eugenol, gallic acid, geranyl acetate, β - <i>N</i> -[γ - <i>l</i> (+)-glutamyl]-4-hydroxy-methylphenylhydrazine, glycyrrhetic acid, <i>p</i> -hydrazinobenzoic acid, isosafrole, kaempferol, <i>d</i> -menthol, nicotine, norharman, pilocarpine, piperidine, protocatechuic acid, rotenone, rutin sulfate, sodium benzoate, turmeric oleoresin, vinblastine

These rodent carcinogens occur in: absinthe, allspice, anise, apple, apricot, banana, basil, beet, broccoli, Brussels sprouts, cabbage, cantaloupe, caraway, cardamom, carrot, cauliflower, celery, cherries, chili pepper, chocolate milk, cinnamon, cloves, cocoa, coffee, collard greens, comfrey herb tea, corn, coriander, currants, dill, eggplant, endive, fennel, garlic, grapefruit, grapes, guava, honey, honeydew melon, horseradish, kale, lemon, lentils, lettuce, licorice, lime, mace, mango, marjoram, mint, mushrooms, mustard, nutmeg, onion, orange, paprika, parsley, parsnip, peach, pear, peas, black pepper, pineapple, plum, potato, radish, raspberries, rhubarb, rosemary, rutabaga, sage, savory, sesame seeds, soybean, star anise, tarragon, tea, thyme, tomato, turmeric, and turnip.

Table 3. Carcinogenicity in rodents of natural chemicals in roasted coffee (51).

Positive: N=19	acetaldehyde, benzaldehyde, benzene, benzofuran, benzo(a)pyrene, caffeic acid, catechol, 1,2,5,6-dibenzanthracene, ethanol, ethylbenzene, formaldehyde, furan, furfural, hydrogen peroxide, hydroquinone, limonene, styrene, toluene, xylene
Not positive: N=8	acrolein, biphenyl, choline, eugenol, nicotinamide, nicotinic acid, phenol, piperidine
Uncertain:	caffeine
Yet to test:	~ 1000 chemicals

Table 4. Proportion of chemicals evaluated as carcinogenic.

Chemicals tested in both rats and mice^a	330/559	(59%)
Naturally-occurring chemicals	73/127	(57%)
Synthetic chemicals	257/432	(59%)
Chemicals tested in rats and/or mice^a		
Chemicals in Carcinogenic Potency Database	668/1275	(52%)
Natural pesticides	35/64	(55%)
Mold toxins	14/23	(61%)
Chemicals in roasted coffee	19/28	(68%)
Innes negative chemicals retested^{a,b}	16/34	(47%)
Physician's Desk Reference (PDR): drugs with reported cancer tests^c	117/241	(49%)
FDA database of drug submissions^d	125/282	(44%)

^a From the Carcinogenic Potency Database (51).

^b The 1969 study by Innes *et al.* (84) is frequently cited as evidence that the proportion of carcinogens is low, as only 9% of 119 chemicals tested (primarily pesticides) were positive. However, these tests, which were only in mice with few animals per group, lacked the power of modern tests. Of the 34 Innes negative chemicals that have been retested using modern protocols: 16 were positive.

^c Davies and Monro (85).

^d Contrera *et al.* (86). 140 drugs are in both the FDA and PDR databases.

Table 5. Ranking Possible Carcinogenic Hazards from Average U.S. Exposures (51). [Chemicals that occur naturally in foods are in bold.] *Daily human exposure:* The calculations assume an average daily dose for a lifetime. *Possible hazard:* The human exposure to a rodent carcinogen is divided by 70 kg to give a mg/kg/day of human exposure, and this dose is given as the percentage of the TD₅₀ in the rodent (mg/kg/day) to calculate the *Human Exposure/Rodent Potency* index (HERP), i.e., 100% means that the human exposure in mg/kg/day is equal to the dose estimated to give 50% of the rodents tumors. TD₅₀ values used in the HERP calculation are averages calculated by taking the harmonic mean of the TD₅₀s of the positive tests in that species from the Carcinogenic Potency Database. Average TD₅₀ values, have been calculated separately for rats and mice, and the more potent value is used for calculating possible hazard. The less potent value is in parentheses.

Possible hazard:	Average daily US exposure	Human dose of rodent carcinogen	Potency TD ₅₀ (mg/kg/day) ^a	
			Rats	Mice
HERP (%)				
140	EDB: workers (high exposure) (before 1977)	Ethylene dibromide, 150 mg	1.52	(7.45)
17	Clofibrate	Clofibrate, 2 g	169	.
14	Phenobarbital, 1 sleeping pill	Phenobarbital, 60 mg	(+)	6.09
6.8	1,3-Butadiene: rubber workers (1978-86)	1,3-Butadiene, 66.0 mg	(261)	13.9
6.1	Tetrachloroethylene: dry cleaners with dry-to-dry units (1980-90) ^b	Tetrachloroethylene, 433 mg	101	(126)
4.0	Formaldehyde: workers	Formaldehyde, 6.1 mg	2.19	(43.9)
2.1	Beer, 257 g	Ethyl alcohol, 13.1 ml	9110	(—)
1.4	Mobile home air (14 hours/day)	Formaldehyde, 2.2 mg	2.19	(43.9)
0.9	Methylene chloride: workers (1940s-80s)	Methylene chloride, 471 mg	724	(918)
0.5	Wine, 28.0 g	Ethyl alcohol, 3.36 ml	9110	(—)
0.4	Conventional home air (14 hours/day)	Formaldehyde, 598 mg	2.19	(43.9)
0.1	Coffee, 13.3 g	Caffeic acid, 23.9 mg	297	(4900)
0.04	Lettuce, 14.9 g	Caffeic acid, 7.90 mg	297	(4900)
0.03	Safrole in spices	Safrole, 1.2 mg	(441)	51.3
0.03	Orange juice, 138 g	d-Limonene, 4.28 mg	204	(—)
0.03	Pepper, black, 446 mg	d-Limonene, 3.57 mg	204	(—)
0.02	Mushroom (<i>Agaricus bisporus</i> 2.55 g)	Mixture of hydrazines, etc. (whole mushroom)	—	20,300
0.02	Apple, 32.0 g	Caffeic acid, 3.40 mg	297	(4900)
0.02	Coffee, 13.3 g	Catechol, 1.33 mg	118	(244)
0.02	Coffee, 13.3 g	Furfural, 2.09 mg	(683)	197
0.009	BHA: daily US avg (1975)	BHA, 4.6 mg	745	(5530)
0.008	Beer (before 1979), 257 g	Dimethylnitrosamine, 726 ng	0.124	(0.189)
0.008	Aflatoxin: daily US avg (1984-89)	Aflatoxin, 18 ng	0.0032	(+)
0.007	Cinnamon, 21.9 mg	Coumarin, 65.0 mg	13.9	(103)
0.006	Coffee, 13.3 g	Hydroquinone, 333 mg	82.8	(225)
0.005	Saccharin: daily US avg (1977)	Saccharin, 7 mg	2140	(—)
0.005	Carrot, 12.1 g	Aniline, 624 mg	194 ^c	(—)
0.004	Potato, 54.9 g	Caffeic acid, 867 mg	297	(4900)
0.004	Celery, 7.95 g	Caffeic acid, 858 mg	297	(4900)
0.004	White bread, 67.6 g	Furfural, 500 mg	(683)	197
0.003	Nutmeg, 27.4 mg	d-Limonene, 466 mg	204	(—)
0.003	Conventional home air (14 hour/day)	Benzene, 155 mg	(169)	77.5
0.002	Carrot, 12.1 g	Caffeic acid, 374 mg	297	(4900)
0.002	Ethylene thiourea: daily US avg (1990)	Ethylene thiourea, 9.51 mg	7.9	(23.5)
0.002	[DDT: daily US avg (before 1972 ban)]	[DDT, 13.8 mg]	(84.7)	12.3
0.001	Plum, 2.00 g	Caffeic acid, 276 mg	297	(4900)

Table 5 contd.

0.001	BHA: daily US avg (1987)	BHA, 700 mg	745	(5530)
0.001	Pear, 3.29 g	Caffeic acid, 240 mg	297	(4900)
0.001	[UDMH: daily US avg (1988)]	[UDMH, 2.82 mg (from Alar)]	(—)	3.96
0.0009	Brown mustard, 68.4 mg	Allyl isothiocyanate, 62.9 mg	96	(—)
0.0008	[DDE: daily US avg (before 1972 ban)]	[DDE, 6.91 mg]	(—)	12.5
0.0007	TCDD: daily US avg (1994)	TCDD, 12.0 pg	0.0000235	(0.000156)
0.0007	Bacon, 11.5 g	Diethylnitrosamine, 11.5 ng	0.0237	(+)
0.0006	Mushroom (<i>Agaricus bisporus</i> 2.55 g)	Glutamyl-<i>p</i>-hydrazinobenzoate, 107 mg	.	277
0.0004	Bacon, 11.5 g	<i>N</i>-Nitrosopyrrolidine, 196 ng	(0.799)	0.679
0.0004	Bacon, 11.5 g	Dimethylnitrosamine, 34.5 ng	0.124	(0.189)
0.0004	[EDB: Daily US avg (before 1984 ban)]	[EDB, 420 ng]	1.52	(7.45)
0.0004	Tap water, 1 liter (1987-92)	Bromodichloromethane, 13 mg	(72.5)	47.7
0.0003	Mango, 1.22 g	<i>d</i>-Limonene, 48.8 mg	204	(—)
0.0003	Beer, 257 g	Furfural, 39.9 mg	(683)	197
0.0003	Tap water, 1 liter (1987-92)	Chloroform, 17 mg	(262)	90.3
0.0003	Carbaryl: daily US avg (1990)	Carbaryl, 2.6 mg	14.1	(—)
0.0002	Celery, 7.95 g	8-Methoxypsoralen, 4.86 mg	32.4	(—)
0.0002	Toxaphene: daily US avg (1990)	Toxaphene, 595 ng	(—)	5.57
0.00009	Mushroom (<i>Agaricus bisporus</i>, 2.55 g)	<i>p</i>-Hydrazinobenzoate, 28 mg	.	454 ^c
0.00008	PCBs: daily US avg (1984-86)	PCBs, 98 ng	1.74	(9.58)
0.00008	DDE/DDT: daily US avg (1990)	DDE, 659 ng	(—)	12.5
0.00007	Parsnip, 54.0 mg	8-Methoxypsoralen, 1.57 mg	32.4	(—)
0.00007	Toast, 67.6 g	Urethane, 811 ng	(41.3)	16.9
0.00006	Hamburger, pan fried, 85 g	PhIP, 176 ng	4.29 ^c	(28.6 ^c)
0.00005	Estragole in spices	Estragole, 1.99 mg	.	51.8
0.00005	Parsley, fresh, 324 mg	8-Methoxypsoralen, 1.17 mg	32.4	(—)
0.00003	Hamburger, pan fried, 85 g	MeIQx, 38.1 ng	1.99	(24.3)
0.00002	Dicofol: daily US avg (1990)	Dicofol, 544 ng	(—)	32.9
0.00001	Cocoa, 3.34 g	α-Methylbenzyl alcohol, 4.3 mg	458	(—)
0.00001	Beer, 257 g	Urethane, 115 ng	(41.3)	16.9
0.000005	Hamburger, pan fried, 85 g	IQ, 6.38 ng	1.89 ^c	(19.6)
0.000001	Lindane: daily US avg (1990)	Lindane, 32 ng	(—)	30.7
0.0000004	PCNB: daily US avg (1990)	PCNB (Quintozene), 19.2 ng	(—)	71.1
0.0000001	Chlorobenzilate: daily US avg (1989)	Chlorobenzilate, 6.4 ng	(—)	93.9
<0.00000001	Chlorothalonil: daily US avg (1990)	Chlorothalonil, <6.4 ng	828 ^d	(—)
0.000000008	Folpet: daily US avg (1990)	Folpet, 12.8 ng	.	2280 ^d
0.000000006	Captan: daily US avg (1990)	Captan, 11.5 ng	2690 ^d	(2730 ^d)

^a“.” = no data in CPDB; (—) = negative in cancer test; (+) = positive cancer test(s) not suitable for calculating a TD₅₀.

^bThis is not an average, but a reasonably large sample (1027 workers).

^cTD₅₀ harmonic mean was estimated for the base chemical from the hydrochloride salt.

^dAdditional data from EPA that is not in the CPDB were used to calculate these TD₅₀ harmonic means.